

Beverly A. H. Buscemi, Ph.D.

State Director

David A. Goodell

*Associate State Director
Operations*

Susan Kreh Beck

*Associate State Director
Policy*

Thomas P. Waring

*Associate State Director
Administration*



3440 Harden Street Ext (29203)
PO Box 4706, Columbia, South Carolina 29240
803/898-9600
Toll Free: 888/DSN-INFO
Website: www.ddsn.sc.gov

COMMISSION

Christine Sharp

Chairperson

Fred Lynn

Vice Chairman

Eva R. Ravenel

Secretary

William O. Danielson

Katherine W. Davis

Katherine Llamas Finley

Harvey E. Shiver

Reference Number: 603-09-DD

Title of Document: Management of Consumers Exposed to Potential Blood borne Pathogens

Date of Issue: March 11, 1996

Effective Date: March 11, 1996

Last Review Date: February 13, 2015

Date of Last Revision: February 13, 2015 **(NO REVISIONS)**

Applicability: DDSN Regional Centers

PURPOSE:

The purpose of this directive is to guide the care of any person residing in a Regional Facility who has an exposure to blood, bloody body fluids or certain body fluids designated as infectious for blood borne pathogens by the Centers for Disease Control and Prevention (CDC). The CDC has designated the following body fluids as infectious for blood borne pathogens: blood, body fluids containing visible blood, semen, vaginal secretions, tissues, cerebrospinal fluid, synovial fluid, pleural fluid, peritoneal fluid, pericardial fluid, and amniotic fluid. [Henceforth, referred to as blood or other potentially infectious materials (OPIM)]. The CDC does not consider the following as infectious unless they contain visible blood: feces, nasal secretions, sputum, sweat, tears, urine, or vomitus.

GENERAL:

Although the potential for Hepatitis B virus (HBV) and Hepatitis C virus (HCV) transmission is greater than that of Human Immunodeficiency Virus (HIV), the transmission of these viruses is similar. These viruses can potentially be transmitted in work sites and living areas through percutaneous inoculation, open wound, non-intact (e.g., chapped, abraded, weeping) skin or mucous membrane contact with blood, bloody body fluids or certain body fluids.

DISTRICT I

P.O. Box 239
Clinton, SC 29325-5328
Phone: (864) 938-3497

Midlands Center - Phone: 803/935-7500
Whitten Center - Phone: 864/833-2733

9995 Miles Jamison Road
Summerville, SC 29485
Phone: 843/832-5576

DISTRICT II

Coastal Center - Phone: 843/873-5750
Pee Dee Center - Phone: 843/664-2600
Saleeby Center - Phone: 843/332-4104

PROCEDURE:

I. TYPES OF EXPOSURE WHICH NEED FURTHER EVALUATION

- A. Percutaneous injury: Blood or OPIM from source person enters another person's skin by:
 - 1. Needle stick;
 - 2. Puncture/cut with a sharp instrument (e.g., razor, knife blade, etc.);
 - 3. Deep scratch: Blood or OPIM must be present on the source person's hands or obviously under fingernails to enter opening in the other person's skin;
 - 4. Bites that break the skin.
- B. Mucous membrane exposure: Blood or OPIM from the source person is spit, splashed, splattered or wiped in the other person's eyes, nose or mouth.
- C. Wound contact: Blood or OPIM from the source person comes into contact with the recipient's skin which has a wound or abrasion less than three (3) days old.
- D. Sexual intercourse.

II. IMMEDIATE FIRST-AID

- A. The injury must immediately or as soon as possible be cleaned.
 - 1. Percutaneous injury: Wash the injury thoroughly with soap and water.
 - 2. Mucous membrane exposure: flush exposed membrane with water for 15 minutes.
 - 3. Wound contact: wash exposed area thoroughly with soap and water.
- B. An injury report must be completed and the incident reported to the unit nurse or the nursing supervisor.

III. DETERMINATION OF EXPOSURE TO A BLOODBORNE PATHOGEN

- A. Once the person has been evaluated by the nurse, the risk of exposure to a blood borne pathogen must be determined as outlined by the Guidelines for Blood Exposure Management at a Glance (Attachment A).


The nurse contacts the attending physician for follow-up using the most recent CDC Guidelines for Managing Occupational Exposures to Blood borne Pathogens (see attachments B, C, D, and E from DDSN Directive 603-05-DD: Policy for Management of Occupational Exposures of Health Care Personnel to Potential Blood Borne Pathogens).

IV. COUNSELING AND FOLLOW UP

- A. A person exposed to HIV, HCV or HBV will be counseled by the appropriate healthcare personnel. The counseling will be conducted according to the person's level of understanding. The person's designated primary contact will be notified of the exposure.
- B. Follow-up testing of the exposed person for HIV will be performed at intervals according to CDC Protocols (usually six weeks, three months, six months, and 12 months) the source person will be tested as needed.



David A. Goodell
Associate State Director-Operations
(Originator)



Beverly A.H. Buscemi, Ph.D.
State Director
(Approved)



Susan Kreh Beck, Ed.S., NCSP
Associate State Director-Policy
(Originator)

To access the following attachments, please see the agency website page "Attachments to Directives" under this directive number.

- | | |
|-----------------|---|
| Attachment A-1: | Consent for HIV/HBV/HCV Testing |
| Attachment A-2: | Guidelines for Blood Exposure Management at a Glance |
| Attachment B: | Blood/Body Fluid Exposure & Testing Summary |
| Attachment C: | Management of Exposure to the Hepatitis B Virus |
| Attachment D: | Management of Exposure to the Hepatitis C Virus |
| Attachment E: | Management of Exposure to the HIV Virus |
| Attachment F: | Situations for which expert consultation for HIV post exposure prophylaxis is advised |